

Social Cooperation Research Program

Project Division of Advanced Biopharmaceutical Science

先進的バイオ医薬品学社会連携研究部門

Project Professor

Kouhei Tsumoto, Ph.D.

Project Associate Professor

Susana de Vega, Ph.D.

特任教授

博士(工学)

津本浩平

特任准教授

博士(生物学)

スサーナ デ ベガ

Various types of antibodies have been approved for therapeutic use and are currently being examined in clinical trials. In the later years, the development of new technologies for the discovery and optimization of high-potency antibodies has led to the ability to design specific and stable antibodies with desired biological properties. Biophysical analyses of therapeutic antibodies, particularly those of protein interaction and stability, are critical for the development of biopharmaceuticals, and an essential step to design next-generation antibodies. The development of analytical methods with quantitative and high-sensitive detection of antigen interaction, protein stability, and biological function of an antibody, therefore, has been challenging for pharmaceutical companies. In this division, we study the biophysical properties of various antibodies to propose a new strategy for the development of highly efficient next-generation antibodies.

1. Experimental modification in thermal stability of oligomers by alanine substitution and site saturation mutagenesis of interfacial residues

Hoya M, Matsunaga R, Nagatoishi S, Tsumoto K.

For certain industrial applications, the stability of protein oligomers is important. In this study, we demonstrated an efficient method to improve the thermal stability of oligomers using the trimeric protein chloramphenicol acetyltransferase (CAT) as the model. We substituted all interfacial residues of CAT with alanine to detect residues critical for oligomer stability. Mutation of six of the forty-nine interfacial residues enhanced oligomer thermal stability. Site saturation mutagenesis was performed on these six residues to optimize the side chains. About 15% of mutations enhanced thermal stability by more than 0.5 °C and most did not disrupt activity of CAT. Certain combinations of mutations further improved thermal stability and resistance against heat treat-

ment. The quadruple mutant, H17V/N34S/F134A/D157C, retained the same activity as the wild-type after heat treatment at 9 °C higher temperature than the wild-type CAT. Furthermore, combinations with only alanine substitutions also improved thermal stability, suggesting the method we developed can be used for rapid modification of industrially important proteins.

2. IL-6 Reduces Spheroid Sizes of Osteophytic Cells Derived from Osteoarthritis Knee Joint via Induction of Apoptosis

Negishi Y, Adili Arepati, Susana de Vega, Momoeda M, Kaneko H, Mehmet Zeynel Cilek, Yoshinaga C, Takafuji K, Otsuka Y, Shimoda M, Negishi-Koga T, Ishijima M, Okada Y

Osteophytes in osteoarthritis (OA) joints contribute to restriction of joint movement, joint pain, and OA progression, but little is known about osteophyte

regulators. Examination of gene expression related to cartilage extracellular matrix, endochondral ossification, and growth factor signaling in articular cartilage and osteophytes obtained from OA knee joints showed that several genes such as COL1A1, VCAN, BGLAP, BMP8B, RUNX2, and SOST were overexpressed in osteophytes compared with articular cartilage. Ratios of mesenchymal stem/progenitor cells, which were characterized by co-expression of CD105 and CD166, were significantly higher in osteophytic cells than articular cells. A three-dimensional culture method for cartilage and osteophyte cells was developed by modification of cultures of self-assembled spheroid cell organoids (spheroids). These spheroids cultured in the media for mesenchymal stem cells containing transforming growth factor- β 3 showed characteristic morphologies and gene expression profiles of articular cartilage and osteophytes, respectively. The effects of IL-1 β , tumor necrosis factor- α , and IL-6 on the spheroids of articular and osteophytic cells were studied. To the best of our knowledge, they provide the first evidence that IL-6 suppresses the spheroid size of osteophytic cells by inducing apoptosis and reducing extracellular matrix molecules. These data show that IL-6 is the suppressor of osteophyte growth and suggest that IL-6 expression and/or activity are implicated in the regulation of osteophyte formation in pathologic joints.

3. Generation of antibodies to an extracellular region of the transporters Glut1/Glut4 by immunization with a designed antigen

Sumikawa T, Nakakido M, Matsunaga R, Kuroda D, Nagatoishi S, Tsumoto K.

Monoclonal antibodies are one of the fastest growing class of drugs. Nevertheless, relatively few biologics target multispinning membrane proteins because of technical challenges. To target relatively small extracellular regions of multiple membrane-spanning proteins, synthetic peptides, which are composed of amino acids corresponding to an extracellular region of a membrane protein, are often utilized in antibody discovery. However, antibodies to these peptides often do not recognize parental membrane proteins. In this study, we designed fusion proteins in which an extracellular helix of the membrane protein glucose transporter 1 (Glut1) was grafted onto the scaffold protein Adhiron. In the initial design, the grafted fragment did not form a helical conformation. Molecular dynamics simulations of full-length Glut1 suggested the importance of intramolecular interactions formed by surrounding residues in the formation of the helical conformation. A fusion protein designed to maintain such intramolecular interactions did form the desired helical conformation in the grafted region. We then immunized an alpaca with the designed fusion protein and obtained VHH (variable region of

heavy-chain antibodies) using the phage display method. The binding of these VHH antibodies to the recombinant Glut1 protein was evaluated by surface plasmon resonance, and their binding to Glut1 on the cell membrane was further validated by flow cytometry. Furthermore, we also succeeded in the generation of a VHH against another integral membrane protein, glucose transporter 4 (Glut4) with the same strategy. These illustrates that our combined biochemical and computational approach can be applied to designing other novel fusion proteins for generating site-specific antibodies.

4. Unveiling the affinity-stability relationship in anti-measles virus antibodies: a computational approach for hotspots prediction

Paul R, Kasahara K, Sasaki J, Pérez JF, Matsunaga R, Hashiguchi T, Kuroda D, Tsumoto K.

Recent years have seen an uptick in the use of computational applications in antibody engineering. These tools have enhanced our ability to predict interactions with antigens and immunogenicity, facilitate humanization, and serve other critical functions. However, several studies highlight the concern of potential trade-offs between antibody affinity and stability in antibody engineering. In this study, we analyzed anti-measles virus antibodies as a case study, to examine the relationship between binding affinity and stability, upon identifying the binding hotspots. We leverage in silico tools like Rosetta and FoldX, along with molecular dynamics (MD) simulations, offering a cost-effective alternative to traditional in vitro mutagenesis. We introduced a pattern in identifying key residues in pairs, shedding light on hotspots identification. Experimental physicochemical analysis validated the predicted key residues by confirming significant decrease in binding affinity for the high-affinity antibodies to measles virus hemagglutinin. Through the nature of the identified pairs, which represented the relative hydropathy of amino acid side chain, a connection was proposed between affinity and stability. The findings of the study enhance our understanding of the interactions between antibody and measles virus hemagglutinin. Moreover, the implications of the observed correlation between binding affinity and stability extend beyond the field of anti-measles virus antibodies, thereby opening doors for advancements in antibody research.

5. THOUSAND-GRAIN WEIGHT 6, which is an IAA-glucose hydrolase, preferentially recognizes the structure of the indole ring

Akabane T, Suzuki N, Ikeda K, Yonezawa T, Nagatoishi S, Matsumura H, Yoshizawa T, Tsuchiya W, Kamino S, Tsumoto K, Ishimaru K, Katoh E, Hirotsu N.

An indole-3-acetic acid (IAA)-glucose hydrolase, THOUSAND-GRAIN WEIGHT 6 (TGW6), negatively regulates the grain weight in rice. TGW6 has been used as a target for breeding increased rice yield. Moreover, the activity of TGW6 has been thought to involve auxin homeostasis, yet the details of this putative TGW6 activity remain unclear. Here, we show the three-dimensional structure and substrate preference of TGW6 using X-ray crystallography, thermal shift assays and fluorine nuclear magnetic resonance (^{19}F NMR). The crystal structure of TGW6 was determined at 2.6 Å resolution and exhibited a six-bladed β -propeller structure. Thermal shift assays revealed that TGW6 preferably interacted with indole compounds among the tested substrates, enzyme products and their analogs. Further analysis using ^{19}F NMR with 1,134 fluorinated fragments emphasized the importance of indole fragments in recognition by TGW6. Finally, docking simulation analyses of the substrate and related fragments in the presence of TGW6 supported the interaction specificity for indole compounds. Herein, we describe the structure and substrate preference of TGW6 for interacting with indole fragments during substrate recognition. Uncovering the molecular details of TGW6 activity will stimulate the use of this enzyme for increasing crop yields and contributes to functional studies of IAA glycoconjugate hydrolases in auxin homeostasis.

6. Structural basis for the recognition of human hemoglobin by the heme-acquisition protein Shr from *Streptococcus pyogenes*

Senoo A, Hoshino M, Shiomi T, Nakakido M, Nagatoishi S, Kuroda D, Nakagawa I, Tame JRH, Caaveiro JMM, Tsumoto K.

In Gram-positive bacteria, sophisticated machineries to acquire the heme group of hemoglobin (Hb) have evolved to extract the precious iron atom contained in it. In the human pathogen *Streptococcus pyogenes*, the Shr protein is a key component of this machinery. Herein we present the crystal structure of hemoglobin-interacting domain 2 (HID2) of Shr bound to Hb. HID2 interacts with both, the protein and heme portions of Hb, explaining the specificity of HID2 for the heme-bound form of Hb, but not its heme-depleted form. Further mutational analysis shows little tolerance of HID2 to interfacial mutations, suggesting that its interaction surface with Hb could be a suitable candidate to develop efficient inhibitors abrogating the binding of Shr to Hb.

7. A high-resolution structural characterization and physicochemical study of how a peptoid binds to an oncoprotein MDM2

Yokomine M, Morimoto J, Fukuda Y, Ueda T, Takeuchi K, Umezawa K, Ago H, Matsuura H, Ueno G,

Senoo A, Nagatoishi S, Tsumoto K, Sando S.

Peptoids are a promising drug modality targeting disease-related proteins, but how a peptoid engages in protein binding is poorly understood. This is primarily due to a lack of high-resolution peptoid-protein complex structures and systematic physicochemical studies. Here, we present the first crystal structure of a peptoid bound to a protein, providing high-resolution structural information about how a peptoid binds to a protein. We previously reported a rigid peptoid, oligo(N-substituted alanine) (oligo-NSA), and developed an oligo-NSA-type peptoid that binds to MDM2. X-ray crystallographic analysis of the peptoid bound to MDM2 showed that the peptoid recognizes the MDM2 surface predominantly through the interaction of the N-substituents, while the main chain acts as a scaffold. Additionally, conformational, thermodynamic, and kinetic analysis of the peptoid and its derivatives with a less rigid main chain revealed that rigidification of the peptoid main chain contributes to improving the protein binding affinity. This improvement is thermodynamically attributed to an increased magnitude of the binding enthalpy change, and kinetically to an increased association rate and decreased dissociation rate. This study provides invaluable insights into the design of protein-targeting peptoids.

8. Thermodynamic and molecular dynamic insights into how fusion influences peptide-tag recognition of an antibody

Miyanabe K, Yamashita T, Tsumoto K.

To understand the effect of protein fusion on the recognition of a peptide-tag by an antibody, we fused a CCR5-derived peptide-tag (pep1) to GFP and investigated its recognition by an anti-pep1 antibody, 4B08. First, to characterize the thermodynamic properties associated with the pep1-4B08 binding, isothermal titration calorimetry experiments were conducted. It was found that pep1 fused to the C-terminus of GFP (GFP-CT) enhanced the enthalpic gain by 2.1 kcal mol⁻¹ and the entropic loss only by 0.9 kcal mol⁻¹, resulting in an 8-fold increase in the binding affinity compared to the unfused pep1. On the other hand, pep1 fused to the N-terminus of GFP (GFP-NT) enhanced the enthalpic gain by 3.0 kcal mol⁻¹ and the entropic loss by 3.2 kcal mol⁻¹, leading to no significant enhancement of the binding affinity. To gain deeper insights, molecular dynamics simulations of GFP-NT, GFP-CT, and pep1 were performed. The results showed that the location of the fusion point sensitively affects the interaction energy, the solvent accessible surface area, and the fluctuation of pep1 in the unbound state, which explains the difference in the experimental thermodynamic properties.

9. Functional insights of Tyr37 in framework region 2 directly contributing to the binding affinities and dissociation kinetics in single-domain VHH antibodies

Yamamoto K, Nagatoishi S, Nakakido M, Kuroda D, Tsumoto K.

Single-domain VHH antibody is regarded as one of the promising antibody classes for therapeutic and diagnostic applications. VHH antibodies have amino acids in framework region 2 that are distinct from those in conventional antibodies, such as the Val-37Phe/Tyr (V37F/Y) substitution. Correlations between the residue type at position 37 and the conformation of the CDR3 in VHH antigen recognition have been previously reported. However, few studies focused on the meaning of harboring two residue types in position 37 of VHH antibodies, and the concrete roles of Y37 have been little to be elucidated. Here, we investigated the functional states of position 37 in co-crystal structures and performed analyses of three model antibodies with either F or Y at position 37. Our analysis indicates that Y at position 37 enhances the dissociation rate, which is highly correlated with drug efficacy. Our findings help to explain the molecular mechanisms that distinguish VHH antibodies from conventional antibodies.

10. Cryo-EM structures elucidate the multiligand receptor nature of megalin

Goto S, Tsutsumi A, Lee Y, Hosojima M, Kabasawa H, Komochi K, Nagatoishi S, Takemoto K, Tsumoto K, Nishizawa T, Kikkawa M, Saito A.

Megalin (low-density lipoprotein receptor-related protein 2) is a giant glycoprotein of about 600 kDa, mediating the endocytosis of more than 60 ligands, including those of proteins, peptides, and drug compounds [S. Goto, M. Hosojima, H. Kabasawa, A. Saito, *Int. J. Biochem. Cell Biol.* 157, 106393 (2023)]. It is expressed predominantly in renal proximal tubule epithelial cells, as well as in the brain, lungs, eyes, inner ear, thyroid gland, and placenta. Megalin is also known to mediate the endocytosis of toxic compounds, particularly those that cause renal and hearing disorders [Y. Hori et al., *J. Am. Soc. Nephrol.* 28, 1783-1791 (2017)]. Genetic megalin deficiency causes Donnai-Barrow syndrome/facio-oculo-acoustico-renal syndrome in humans. However, it is not known how megalin interacts with such a wide variety of ligands and plays pathological roles in various organs. In this study, we elucidated the dimeric architecture of megalin, purified from rat kidneys, using cryoelectron microscopy. The maps revealed the densities of endogenous ligands bound to various regions throughout the dimer, elucidating the multiligand receptor nature of megalin. We also determined the

structure of megalin in complex with receptor-associated protein, a molecular chaperone for megalin. The results will facilitate further studies on the pathophysiology of megalin-dependent multiligand endocytic pathways in multiple organs and will also be useful for the development of megalin-targeted drugs for renal and hearing disorders, Alzheimer's disease [B. V. Zlokovic et al., *Proc. Natl. Acad. Sci. U.S.A.* 93, 4229-4234 (1996)], and other illnesses.

11. Next-Generation Anti-TNF α Agents: The Example of Ozoralizumab

Tsumoto K, Takeuchi T.

Biologic therapy involving anti-tumor necrosis factor- α (anti-TNF α) agents has fundamentally changed the management of patients with immune-mediated inflammatory diseases, including rheumatoid arthritis, thus benefiting many patients. Nevertheless, the inability of some patients to achieve low disease activity or clinical remission remains a major concern. To address such concerns, next-generation anti-TNF α agents that differ from the immunoglobulin G-format anti-TNF α agents that have been used to date are being developed using antibody-engineering technology. Their unique design employing novel molecular characteristics affords several advantages, such as early improvement of clinical symptoms, optimization of drug bioavailability, enhancement of tissue penetration, and a reduction in side effects. This holds promise for a new paradigm shift in biologic therapy via the use of next-generation anti-TNF α agents. Ozoralizumab, a next-generation anti-TNF α agent that was recently approved in Japan, comprises a variable region heavy-chain format. It has a completely different structure from conventional therapeutic antibodies, such as a small molecular size, an albumin-binding module, and a unique format that produces an avidity effect. Ozoralizumab exhibited rapid biodistribution into joints, provided attenuation of Fc γ receptor-mediated inflammatory responses, and had a high binding affinity to TNF α in non-clinical studies. In clinical trials, ozoralizumab yielded an early improvement in clinical symptoms, a sustained efficacy for up to 52 weeks, and an acceptable tolerability in patients with rheumatoid arthritis. This review focuses on the results of pre-clinical and clinical trials for ozoralizumab and outlines the progress in next-generation antibody development.

12. Characterization of a novel format scFv \times VHH single-chain biparatopic antibody against metal binding protein MtsA

Asano R, Takeuchi M, Nakakido M, Ito S, Aikawa C, Yokoyama T, Senoo A, Ueno K, Nagatoishi S, Tanaka Y, Nakagawa I, Tsumoto K.

Biparatopic antibodies (bpAbs) are engineered antibodies that bind to multiple different epitopes within the same antigens. bpAbs comprise diverse formats, including fragment-based formats, and choosing the appropriate molecular format for a desired function against a target molecule is a challenging task. Moreover, optimizing the design of constructs requires selecting appropriate antibody modalities and adjusting linker length for individual bpAbs. Therefore, it is crucial to understand the characteristics of bpAbs at the molecular level. In this study, we first obtained single-chain variable fragments and camelid heavy-chain variable domains targeting distinct epitopes of the metal binding protein MtsA and then developed a novel format single-chain bpAb connecting these fragment antibodies with various linkers. The physicochemical properties, binding activities, complex formation states with antigen, and functions of the bpAb were analyzed using multiple approaches. Notably, we found that the assembly state of the complexes was controlled by a linker and that longer linkers tended to form more compact complexes. These observations provide detailed molecular information that should be considered in the design of bpAbs.

13. Crystal structures of human CD40 in complex with monoclonal antibodies dacetuzumab and bleselumab

Asano R, Nakakido M, Pérez JF, Ise T, Caaveiro JMM, Nagata S, Tsumoto K.

CD40 is a member of the tumor necrosis factor receptor superfamily, and it is widely expressed on immune and non-immune cell types. The interaction between CD40 and the CD40 ligand (CD40L) plays an essential function in signaling, and the CD40/CD40L complex works as an immune checkpoint molecule. CD40 has become a therapeutic target, and a variety of agonistic/antagonistic anti-CD40 monoclonal antibodies (mAbs) have been developed. To better understand the mode of action of anti-CD40 mAbs, we determined the X-ray crystal structures of dacetuzumab (agonist) and bleselumab (antagonist) in complex with the extracellular domain of human CD40, respectively. The structure reveals that dacetuzumab binds to CD40 on the top of cysteine-rich domain 1 (CRD1), which is the domain most distant from the cell surface, and it does not compete with CD40L binding. The binding interface of bleselumab spread between CRD2 and CRD1, overlapping with the binding surface of the ligand. Our results offer important insights for future structural and functional studies of CD40 and provide clues to understanding the mechanism of biological response. These data can be applied to developing new strategies for designing antibodies with more therapeutic efficacy.

14. High-throughput system for the thermostability analysis of proteins

Ito S, Matsunaga R, Nakakido M, Komura D, Katoh H, Ishikawa S, Tsumoto K.

Thermal stability of proteins is a primary metric for evaluating their physical properties. Although researchers attempted to predict it using machine learning frameworks, their performance has been dependent on the quality and quantity of published data. This is due to the technical limitation that thermodynamic characterization of protein denaturation by fluorescence or calorimetry in a high-throughput manner has been challenging. Obtaining a melting curve that derives solely from the target protein requires laborious purification, making it far from practical to prepare a hundred or more samples in a single workflow. Here, we aimed to overcome this throughput limitation by leveraging the high protein secretion efficacy of *Brevibacillus* and consecutive treatment with plate-scale purification methodologies. By handling the entire process of expression, purification, and analysis on a per-plate basis, we enabled the direct observation of protein denaturation in 384 samples within 4 days. To demonstrate a practical application of the system, we conducted a comprehensive analysis of 186 single mutants of a single-chain variable fragment of nivolumab, harvesting the melting temperature (T_m) ranging from -9.3 up to $+10.8^\circ\text{C}$ compared to the wild-type sequence. Our findings will allow for data-driven stabilization in protein design and streamlining the rational approaches.

15. Impact of single-residue mutations on protein thermal stability: The case of threonine 83 of BC2L-CN lectin

Hoya M, Matsunaga R, Nagatoishi S, Ide T, Kuroda D, Tsumoto K.

The thermal stability of trimeric lectin BC2L-CN was investigated and found to be considerably altered when mutating residue 83, originally a threonine, located at the fucose-binding loop. Mutants were analyzed using differential scanning calorimetry and isothermal microcalorimetry. Although most mutations decreased the affinity of the protein for oligosaccharide H type 1, six mutations increased the melting temperature (T_m) by $>5^\circ\text{C}$; one mutation, T83P, increased the T_m value by 18.2°C ($T83P$, $T_m = 96.3^\circ\text{C}$). In molecular dynamic simulations, the investigated thermostable mutants, T83P, T83A, and T83S, had decreased fluctuations in the loop containing residue 83. In the T83S mutation, the side-chain hydroxyl group of serine formed a hydrogen bond with a nearby residue, suggesting that the restricted movement of the side-chain resulted in fewer fluctuations and enhanced thermal stability. Residue 83 is located

at the interface and near the upstream end of the equivalent loop in a different protomer; therefore, fluctuations by this residue likely propagate throughout the loop. Our study of the dramatic change in thermal stability by a single amino acid mutation provides useful insights into the rational design of protein structures, especially the structures of oligomeric proteins.

16. PDZD8-FKBP8 tethering complex at ER-mitochondria contact sites regulates mitochondrial complexity

Nakamura K, Aoyama-Ishiwatari S, Nagao T, Paaran M, Obara CJ, Sakurai-Saito Y, Johnston J, Du Y, Suga S, Tsuboi M, Nakakido M, Tsumoto K, Kishi Y, Gotoh Y, Kwak C, Rhee HW, Seo JK, Kosako H, Potter C, Carragher B, Lippincott-Schwartz J, Polleux F, Hirabayashi Y.

Mitochondria-ER membrane contact sites (MERCS) represent a fundamental ultrastructural feature underlying unique biochemistry and physiology in eukaryotic cells. The ER protein PDZD8 is required for the formation of MERCS in many cell types, however, its tethering partner on the outer mitochondrial membrane (OMM) is currently unknown. Here we identified the OMM protein FKBP8 as the tethering partner of PDZD8 using a combination of unbiased proximity proteomics, CRISPR-Cas9 endogenous protein tagging, Cryo-Electron Microscopy (Cryo-EM) tomography, and correlative light-EM (CLEM). Single molecule tracking revealed highly dynamic diffusion properties of PDZD8 along the ER membrane with significant pauses and capture at MERCS. Overexpression of FKBP8 was sufficient to narrow the ER-OMM distance, whereas independent versus combined deletions of these two proteins demonstrated their interdependence for MERCS formation. Furthermore, PDZD8 enhances mitochondrial complexity in a FKBP8-dependent manner. Our results identify a novel ER-mitochondria tethering complex that regulates mitochondrial morphology in mammalian cells.

17. Development of novel humanized VHH synthetic libraries based on physicochemical analyses

Nakakido M, Kinoshita S, Tsumoto K.

Due to the high affinity and specificity of antibodies toward antigens, various antibody-based applications have been developed. Recently, variable antigen-binding domains of heavy-chain antibodies (VHH) have become an attractive alternative to conventional fragment antibodies due to their unique molecular characteristics. As an antibody-generating strategy, synthetic VHH libraries (including human-

ized VHH libraries) have been developed using distinct strategies to constrain the diversity of amino acid sequences. In this study, we designed and constructed several novel synthetic humanized VHH libraries based on biophysical analyses conducted using the complementarity determining region-grafting method and comprehensive sequence analyses of VHHs deposited in the protein data bank. We obtained VHHs from the libraries, and hit clones exhibited considerable thermal stability. We also found that VHHs from distinct libraries tended to have different epitopes. Based on our results, we propose a strategy for generating humanized VHHs with distinct epitopes toward various antigens by utilizing our library combinations.

18. Systematic Preparation of a 66-IgG Library with Symmetric and Asymmetric Homogeneous Glycans and Their Functional Evaluation

Manabe S, Iwamoto S, Nagatoishi S, Hoshino A, Mitani A, Sumiyoshi W, Kinoshita T, Yamaguchi Y, Tsumoto K.

Immunoglobulin G (IgG) antibodies possess a conserved N-glycosylation site in the Fc domain. In FcγRIIIa affinity column chromatography, unglycosylated, hemiglycosylated, and fully glycosylated IgG retention times differ considerably. Using retention-time differences, 66 different trastuzumab antibodies with symmetric and asymmetric homogeneous glycans were prepared systematically, substantially expanding the scope of IgGs with homogeneous glycans. Using the prepared trastuzumab with homogeneous glycans, thermal stability and antibody-dependent cellular cytotoxicity were investigated. In some glycan series, a directly proportional relationship was observed between the thermal unfolding temperature (T_m) and the calorimetric unfolding heat (ΔH_{cal}). Antibody function could be deduced from the combination of a pair of glycans in an intact form. Controlling glycan structure through the combination of a pair of glycans permits the precise tuning of stability and effector functions of IgG. Overall, our technology can be used to investigate the effects of glycans on antibody functions.

19. Triphenylphosphonium-modified cationomers enhance in vivo mRNA delivery through stabilized polyion complexation

Norimatsu J, Mizuno HL, Watanabe T, Obara T, Nakakido M, Tsumoto K, Cabral H, Kuroda D, Anraku Y.

Nanocarriers based on cationic materials play a central role in the success of mRNA-based therapies. Traditionally, amine-bearing lipids and polymers have been successfully employed for creating mR-

NA-loaded nanocarriers, though they still present challenges, such as physical and biological instability, limiting both delivery efficiency and therapeutic potential. Non-amine cations could be a promising avenue in addressing these limitations. However, such alternatives remain notably underexplored. Herein, we introduced triphenylphosphonium (TPP) as an alternative cationic moiety for mRNA delivery, leveraging its advantageous properties for nucleic acid complexation. Through the modification of amine-bearing cationomers, we replaced traditional amine-based counterparts with TPP to create innovative polymeric micelles as mRNA nanocarriers. A comprehensive analysis, encompassing physicochemical, thermodynamic, and computational approaches, revealed that the TPP substitution significantly influenced polymer self-assembly, mRNA binding, and the overall stability of mRNA-loaded polymeric micelles. Upon intravenous injection, TPP-bearing micelles demonstrated a remarkable increase in mRNA bioavailability, facilitating efficient protein production in solid tumors. These findings provide a compelling rationale for substituting amines with TPP, emphasizing their potential for advancing mRNA therapeutics.

20. Unveiling the structural mechanisms behind high affinity and selectivity in phosphorylated epitope-specific rabbit antibodies

Kasahara K, Kawade R, Nakakido M, Matsunaga R, Akiba H, Entzminger KC, Maruyama T, Okumura SCJ, Caaveiro JMM, Kuroda D, Tsumoto K.

Protein phosphorylation is a crucial process in various cellular functions, and its irregularities have been implicated in several diseases, including cancer. Antibodies are commonly employed to detect protein phosphorylation in research. However, unlike the extensive studies on recognition mechanisms of the phosphate group by proteins such as kinases and phosphatases, only a few studies have explored antibody mechanisms. In this study, we produced and characterized two rabbit monoclonal antibodies that recognize a monophosphorylated Akt peptide. Through crystallography, thermodynamic mutational analyses, and molecular dynamics simulations, we investigated the unique recognition mechanism that enables higher binding affinity and selectivity of the antibodies compared to other generic proteins with lower binding affinity to phosphorylated epitopes. Our results demonstrate that molecular dynamics simulations provide novel insights into the dynamic aspects of molecular recognition of posttranslational modifications by proteins beyond static crystal structures, highlighting how specific atomic level interactions drive the exceptional affinity and selectivity of antibodies.

21. Structural basis for the ligand promiscuity of the hydroxamate siderophore binding protein FtsB from *Streptococcus pyogenes*

Fernandez-Perez J, Senoo A, Caaveiro JMM, Nakakido M, de Vega S, Nakagawa I, Tsumoto K.

Pathogenic bacteria must secure the uptake of nutritional metals such as iron for their growth, making their import systems attractive targets for the development of new antimicrobial modalities. In the pathogenic bacterium *Streptococcus pyogenes*, the iron uptake system FtsABCD transports iron encapsulated by siderophores of the hydroxamate class. However, the inability of *S. pyogenes* to produce these metabolites makes the biological and clinical relevance of this route unresolved. Herein, we demonstrated that the periplasmic binding protein FtsB recognizes not only the hydroxamate siderophore ferrichrome, as previously documented, but also ferrioxamine E (FOE), ferrioxamine B (FOB), and bisucaberin (BIS), each of them with high affinity (nM level). Up to seven aromatic residues in the binding pocket accommodate the variable backbones of the different siderophores through CH- π interactions, explaining ligand promiscuity. Collectively, our observations revealed how *S. pyogenes* exploits the diverse xenosiderophores produced by other microorganisms as iron sources to secure this precious nutrient.

22. Malaria parasites require a divergent heme oxygenase for apicoplast gene expression and biogenesis

Blackwell AM, Jami-Alahmadi Y, Nasamu AS, Kudo S, Senoo A, Slam C, Tsumoto K, Wohlschlegel JA, Manuel Martinez Caaveiro J, Goldberg DE, Sigala PA.

Malaria parasites have evolved unusual metabolic adaptations that specialize them for growth within heme-rich human erythrocytes. During blood-stage infection, *Plasmodium falciparum* parasites internalize and digest abundant host hemoglobin within the digestive vacuole. This massive catabolic process generates copious free heme, most of which is biomineralized into inert hemozoin. Parasites also express a divergent heme oxygenase (HO)-like protein (PfHO) that lacks key active-site residues and has lost canonical HO activity. The cellular role of this unusual protein that underpins its retention by parasites has been unknown. To unravel PfHO function, we first determined a 2.8 Å-resolution X-ray structure that revealed a highly α -helical fold indicative of distant HO homology. Localization studies unveiled PfHO targeting to the apicoplast organelle, where it is imported and undergoes N-terminal processing but retains most of the electropositive transit peptide. We observed that conditional knockdown of PfHO was le-

thal to parasites, which died from defective apicoplast biogenesis and impaired isoprenoid-precursor synthesis. Complementation and molecular-interaction studies revealed an essential role for the electropositive N-terminus of PfHO, which selectively associates with the apicoplast genome and enzymes involved in nucleic acid metabolism and gene expression. PfHO knockdown resulted in a specific deficiency in levels of apicoplast-encoded RNA but not DNA. These studies reveal an essential function for PfHO in apicoplast maintenance and suggest that *Plasmodium* repurposed the conserved HO scaffold from its canonical heme-degrading function in the ancestral chloroplast to fulfill a critical adaptive role in organelle gene expression.

23. Disrupted odontoblast differentiation and dentin dysplasia in Epiprofin-deficient mice

Jiménez-Rojo L, de Vega S, Ibarretxe G, Nakamura T, Unda FJ.

Tooth formation is a process tightly regulated by reciprocal interactions between epithelial and mesenchymal tissues. These epithelial-mesenchyme interactions regulate the expression of target genes via transcription factors. Among the regulatory elements governing this process, Epiprofin/Sp6 is a zinc finger transcription factor which is expressed in the embryonic dental epithelium and in differentiating pre-odontoblasts. Epiprofin knockout (Epfⁿ-/-) mice present severe dental abnormalities, such as supernumerary teeth and enamel hypoplasia. Here, we describe dentin defects in molars and incisors of Epfⁿ-/- mice. We observed that in the absence of Epfⁿ, markers of early odontoblast differentiation, such as alkaline phosphatase activity, Dsp/Dpp expression, and Collagen Type I deposition, are downregulated. In addition, the expression of tight and gap junction proteins was severely impaired in the predontoblastic cell layer of developing Epfⁿ-/- molars. Altogether, our data shows that Epfⁿ is crucial for the proper differentiation of dental mesenchymal cells towards functional odontoblasts and subsequent dentin-matrix deposition.

24. Specific recognition mechanism of an antibody to sulfated tyrosine and its potential use in biological research

Ujiie K, Nakakido M, Kinoshita S, Jose Caaveiro M M, Entzminger, C J Okumura, Maruyama, Miyauchi k, Matano T, Tsumoto K

Post-translational modification of proteins is a crucial biological reaction that regulates protein func-

tions by altering molecular properties. The specific detection of such modifications in proteins has made significant contributions to molecular biology research and holds potential for future drug development applications. In HIV research, for example, tyrosine sulfation at the N-terminus of C-C chemokine receptor type 5 (CCR5) is considered to significantly enhance HIV infection efficiency. However, antibodies specific to sulfated CCR5 still need to be developed. In this study, we successfully generated an antibody that specifically recognized the sulfated N-terminal peptide of CCR5 through rabbit immunization and panning via phage display using a CCR5 N-terminal peptide containing sulfate modification. We used various physicochemical methods in combination with molecular dynamics simulation to screen for residues that could be involved in recognition of the sulfated peptide by this antibody. We also confirmed that this antibody recognized the sulfated full-length CCR5 on the cell surface, which suggested it should be useful as a research tool that could lead to the development of novel therapeutics. Although the antibody binding did not inhibit HIV infection, it could be also described as sulfation site-specific binding, beyond sulfation-specific binding.

25. The pericellular function of Fibulin-7 in the adhesion of oligodendrocyte lineage cells to neuronal axons during CNS myelination

Yamada M, Sasaki B, Yamada N, Hayashi C, Tsumoto K, de Vega S, Suzuki N

Myelin is an electrical insulator that enables saltatory nerve conduction and is essential for proper functioning of the central nervous system (CNS). It is formed by oligodendrocytes (OLs) in the CNS, and during OL development various molecules, including extracellular matrix (ECM) proteins, regulate OL differentiation and myelination; however, the role of ECM proteins in these processes is not well understood. Our present work is centered on the analyses of the expression and function of fibulin-7 (Fbln7), an ECM protein of the fibulin family, in OL differentiation. In the expression analysis of Fbln7 in the CNS, we found that it was expressed at early postnatal stage and localized in the processes of OL precursor cells (OPCs), in the inner region of myelin, and in axons. The functional analysis using recombinant Fbln7 protein (rFbln7) revealed that rFbln7 promoted OPC attachment activity via β 1 integrin and heparan sulfate receptors. Further, rFbln7 induced the adhesion to neurites and the differentiation of OLs. Altogether, our results show that Fbln7 promotes the adhesion between OLs and axons and OL differentiation.

Publications

- Hoya M, Matsunaga R, Nagatoishi S, Tsumoto K. Experimental modification in thermal stability of oligomers by alanine substitution and site saturation mutagenesis of interfacial residues. *Biochem Biophys Res Commun.* 691. 149316.2024
- Negishi Y, Adili Arepati, Susana de Vega, Momoe-da M, Kaneko H, Mehmet Zeynel Cilek, Yoshinaga C, Takafuji K, Otsuka Y, Shimoda M, Negishi-Koga T, Ishijima M, Okada Y. IL-6 Reduces Spheroid Sizes of Osteophytic Cells Derived from Osteoarthritis Knee Joint via Induction of Apoptosis. *Am J Pathology.* 194(1). 135-149.2024
- Sumikawa T, Nakakido M, Matsunaga R, Kuroda D, Nagatoishi S, Tsumoto K. Generation of antibodies to an extracellular region of the transporters Glut1/Glut4 by immunization with a designed antigen. *J Biol Chem.* 300. 105640.2024
- Unveiling the affinity-stability relationship in anti-measles virus antibodies: a computational approach for hotspots prediction. Paul R, Kasahara K, Sasaki J, Pérez JF, Matsunaga R, Hashiguchi T, Kuroda D, Tsumoto K. *Front Mol Biosci.* 10.2024
- Akabane T, Suzuki N, Ikeda K, Yonezawa T, Nagatoishi S, Matsumura H, Yoshizawa T, Tsuchiya W, Kamino S, Tsumoto K, Ishimaru K, Katoh E, Hirotsu N. THOUSAND-GRAIN WEIGHT 6, which is an IAA-glucose hydrolase, preferentially recognizes the structure of the indole ring. *Sci Rep.* 14(1). 6778.2024
- Senoo A, Hoshino M, Shiomi T, Nakakido M, Nagatoishi S, Kuroda D, Nakagawa I, Tame JRH, Caaveiro JMM, Tsumoto K. Structural basis for the recognition of human hemoglobin by the heme-acquisition protein Shr from *Streptococcus pyogenes*. *Sci Rep.* 14(1). 5374
- Yokomine M, Morimoto J, Fukuda Y, Ueda T, Takeuchi K, Umezawa K, Ago H, Matsuura H, Ueno G, Senoo A, Nagatoishi S, Tsumoto K, Sando S. A high-resolution structural characterization and physicochemical study of how a peptoid binds to an oncoprotein MDM2. *Chem Sci.* 15(19). 7051-7060.2024
- Miyanabe K, Yamashita T, Tsumoto K. Thermodynamic and molecular dynamic insights into how fusion influences peptide-tag recognition of an antibody. *Sci Rep.* 14(1). 8685.2024
- Yamamoto K, Nagatoishi S, Nakakido M, Kuroda D, Tsumoto K. Functional insights of Tyr37 in framework region 2 directly contributing to the binding affinities and dissociation kinetics in single-domain VHH antibodies. *Biochem Biophys Res Commun.* 709. 149839.2024
- Goto S, Tsutsumi A, Lee Y, Hosojima M, Kabasawa H, Komochi K, Nagatoishi S, Takemoto K, Tsumoto K, Nishizawa T, Kikkawa M, Saito A. Cryo-EM structures elucidate the multiligand receptor nature of megalin. *Proc Natl Acad Sci U S A.* 121(22). e2318859121.2024
- Tsumoto K, Takeuchi T. Next-Generation Anti-TNF α Agents: The Example of Ozoralizumab. *BioDrugs.* 38(3). 341-351.2024
- Asano R, Takeuchi M, Nakakido M, Ito S, Aikawa C, Yokoyama T, Senoo A, Ueno G, Nagatoishi S, Tanaka Y, Nakagawa I, Tsumoto K. Characterization of a novel format scFv \times VHH single-chain biparatopic antibody against metal binding protein MtsA. *Protein Sci.* 33(6). 5017.2024
- Asano R, Nakakido M, Pérez JF, Ise T, Caaveiro JMM, Nagata S, Tsumoto K. Crystal structures of human CD40 in complex with monoclonal antibodies dacetuzumab and bleselumab. *Biochem Biophys Res Commun.* 714. 149969.2024
- Ito S, Matsunaga R, Nakakido M, Komura D, Katoh H, Ishikawa S, Tsumoto K. High-throughput system for the thermostability analysis of proteins. *Protein Sci.* 33(6). e5029.2054
- Hoya M, Matsunaga R, Nagatoishi S, Ide T, Kuroda D, Tsumoto K. Impact of single-residue mutations on protein thermal stability: The case of threonine 83 of BC2L-CN lectin. *Int J Biol Macromol.* 272(Pt 1). 132682.2024
- Nakamura K, Aoyama-Ishiwatari S, Nagao T, Paaran M, Obara CJ, Sakurai-Saito Y, Johnston J, Du Y, Suga S, Tsuboi M, Nakakido M, Tsumoto K, Kishi Y, Gotoh Y, Kwak C, Rhee HW, Seo JK, Kosako H, Potter C, Carragher B, Lippincott-Schwartz J, Polleux F, Hirabayashi Y. PDZD8-FKBP8 tethering complex at ER-mitochondria contact sites regulates mitochondrial complexity. *bioRxiv [Preprint]*. 2023/08/22. 554218.2024
- Nakakido M, Kinoshita S, Tsumoto K. Development of novel humanized VHH synthetic libraries based on physicochemical analyses. *Sci Rep.* 14(1). 19533.2024
- Manabe S, Iwamoto S, Nagatoishi S, Hoshino A, Mitani A, Sumiyoshi W, Kinoshita T, Yamaguchi Y, Tsumoto K. Systematic Preparation of a 66-IgG Library with Symmetric and Asymmetric Homogeneous Glycans and Their Functional Evaluation. *J Am Chem Soc.* 146(33). 23426-23436.2024
- Norimatsu J, Mizuno HL, Watanabe T, Obara T, Nakakido M, Tsumoto K, Cabral H, Kuroda D, Anraku Y. Triphenylphosphonium-modified cationomers enhance in vivo mRNA delivery through stabilized polyion complexation. *Mater Horiz.* 11(19). 4711-4721.2024
- Kasahara K, Kawade R, Nakakido M, Matsunaga R, Akiba H, Entzminger KC, Maruyama T, Okumura SCJ, Caaveiro JMM, Kuroda D, Tsumoto K. Unveiling the structural mechanisms behind high affinity and selectivity in phosphorylated epitope-specific rabbit antibodies. *J Biol Chem.* 300(12). 107989.2024
- Fernandez-Perez J, Senoo A, Caaveiro JMM, Na-

- kakido M, de Vega S, Nakagawa I, Tsumoto K. Structural basis for the ligand promiscuity of the hydroxamate siderophore binding protein FtsB from *Streptococcus pyogenes*. *Structure*. 32(12). 2410-2421.2024
22. Blackwell AM, Jami-Alahmadi Y, Nasamu AS, Kudo S, Senoo A, Slam C, Tsumoto K, Wohlschlegel JA, Manuel Martinez Caaveiro J, Goldberg DE, Sigala PA. Malaria parasites require a divergent heme oxygenase for apicoplast gene expression and biogenesis. *Elife*. 13. RP100256.2024
23. Jiménez-Rojo L, de Vega S, Ibarretxe G, Nakamura T, Unda FJ. Disrupted odontoblast differentiation and dentin dysplasia in Epiprofin-deficient mice . *Int. J Dev. Biol* . 68 (1) . 19-24 .2024
24. Ujiie K, Nakakido M, Kinoshita S, Jose Caaveiro M M, Entzminger, C J Okumura, Maruyama, Miyauchi k, Matano T, Tsumoto K. Specific recognition mechanism of an antibody to sulfated tyrosine and its potential use in biological research. *J Biol Chem*. 108176.2025
25. Yamada M, Sasaki B, Yamada N, Hayashi C, Tsumoto K, de Vega S, Suzuki N. The pericellular function of Fibulin-7 in the adhesion of oligodendrocyte lineage cells to neuronal axons during CNS myelination. *Biochemical and Biophysical Research Communications*.748.151271.2025