

IMSUT Hospital

Department of Radiology

放射線科

Associate Professor	Hiroyuki Akai, M.D., D.M.Sc.
Senior Assistant Professor	Toshihiro Furuta, M.D., D.M.Sc.
Assistant Professor	Haruomi Yamaguchi, M.D., D.M.Sc.
Project Assistant Professor	Shimpei Kato, M.D., D.M.Sc.

准教授	博士(医学)	赤井宏行
講師	博士(医学)	古田寿宏
助教	博士(医学)	山口晴臣
特任助教	博士(医学)	加藤伸平

Department of Radiological Technology

放射線部

Associate Professor	Hiroyuki Akai, M.D., D.M.Sc.
Head Radiologic Technologist	Kenji Ino, RT

准教授	博士(医学)	赤井宏行
放射線技師長		井野賢二

The Department of Radiology undertakes radiology service at IMSUT hospital. Our expertise includes general diagnostic radiology, neuroradiology, clinical nuclear medicine, and radiation therapy. Board-certified radiologists at the Department of Radiology conduct all examinations of CT, MRI, and nuclear medicine. Radiological reports are made by the radiologists. In addition, several clinical studies are being conducted in collaboration with other departments or institutions. We also investigate the technical aspects of molecular imaging with intact small animals for its application to preclinical studies using an optical imaging system and MRI.

The Department of Radiological Technology constitutes the hospital radiology service together with the Department of Radiology. Plain radiography, dual-energy X-ray absorptiometry, and barium studies are also available at the Department of Radiological Technology, other than CT, MRI, and radioisotope examinations. More than 10,000 patients visit our department every year. Radiologic technologists at the department make an effort to provide high-quality medical images in daily practice as well as to reasonably reduce radiation exposure of a patient during the examination.

Feasibility study of direct CT lymphangiography in mice: comparison with interstitial CT/MR lymphangiography.

Sugawara H, Furuta T, Sumiyoshi A¹, Iiyama M¹, Kamitani M, Suzuki A², Murakami A², Abe O³, Aoki I¹, Akai H.

¹ Functional and Molecular Imaging Group, Department of Molecular Imaging and Theranostics, Institute for Quantum Medical Science, National Institutes for Quantum Science and Technology, ² Department of Diagnostic Radiology, Kanto Rosai Hospital, ³ Department of Radiology, Graduate School of Medicine, University of Tokyo

In the present study, we aimed to establish a CT lymphangiography method in mice via direct lymph node puncture. We injected healthy mice ($n = 8$) with 50 μl of water-soluble iodine contrast agent (iomeprol; iodine concentration, 350 mg/mL) subcutaneously into the left-rear foot pad (interstitial injection) and 20 μl of the same contrast agent directly into the popliteal lymph node (direct puncture) 2 days later. Additionally, we performed interstitial MR lymphangiography on eight mice as a control group. We calculated the contrast ratio for each lymph node and visually assessed the depiction of lymph nodes and lymphatic vessels on a three-point scale. As a result, the contrast ratios of 2-min post-injection images of sacral and lumbar-aortic lymph nodes were 20.7 ± 16.6 (average \pm standard deviation) and 17.1 ± 12.0 in the direct puncture group, which were significantly higher than those detected in the CT or MR interstitial lymphangiography groups (average, 1.8–3.6; $p = 0.008$ – 0.019). The visual assessment scores for sacral lymph nodes, lumbar-aortic lymph nodes, and cisterna chyli were significantly better in the direct puncture group than in the CT interstitial injection group ($p = 0.036$, 0.009 and 0.001 , respectively). The lymphatic vessels between these structures were significantly better scored in direct puncture group than in the CT or MR interstitial lymphangiography groups at 2 min after injection (all $p \leq 0.05$). To sum up, in CT lymphangiography in mice, the direct lymph node puncture provides a better delineation of the lymphatic pathways than the CT/MR interstitial injection method, which we believe that direct CT lymphangiography will be the first choice for the evaluation of lymphatic pathways in mice.

Early detection of hypervascularization in hepatocellular carcinoma (≤ 2 cm) on hepatic arterial phase with virtual monochromatic imaging: Comparison with low-tube voltage CT.

Yamaguchi H, Ichikawa T⁴, Morisaka H⁵, Akai H, Izuka K⁶, Ueno T⁶, Abe O³, Tsushima Y⁴.

⁴ Diagnostic Radiology and Nuclear Medicine Department, Gunma University Graduate School of Medicine, ⁵ Department of Radiology, University of Yamanashi, ⁶ Department of Gastroenterology, Isesazaki Municipal Hospital

This study aims to assess the diagnostic value of virtual monochromatic image (VMI) at low keV energy for early detection of small hepatocellular carcinoma (HCC) in hepatic arterial phase compared with low-tube voltage (80 kVp) CT generated from dual-energy CT (DE-CT). A total of 107 patients with 114 hypervascular HCCs (≤ 2 cm) underwent DE-CT, 140 kVp, blended 120 kVp, and 80 kVp images were generated, as well as 40 and 50 keV. CT numbers of HCCs and the standard deviation as image noise on psoas muscle were measured. The contrast-to-noise ratios (CNR) of HCC were compared among all techniques. Overall image quality and sensitivity for detecting HCC hypervascularity were qualitatively assessed by three readers. The mean CT numbers, CNR, and image noise were highest at 40 keV followed by 50 keV, 80 kVp, blended 120 kVp, and 140 kVp. Significant differences were found in all evaluating endpoints except for mean image noise of 50 keV and 80 kVp. Image quality of 40 keV was the lowest, but still it was considered acceptable for diagnostic purposes. The mean sensitivity for detecting lesion hypervascularity with 40 keV (92%) and 50 keV (84%) was higher than those with 80 kVp (56%). Low keV energy images were superior to 80 kVp in detecting hypervascularization of early HCC.

Publications

1. Tajima T, Akai H, Yasaka K, Kunimatsu A, Yamashita Y, Akahane M, Yoshioka N, Abe O, Ohtomo K, Kiryu S. Usefulness of deep learning-based noise reduction for 1.5 T MRI brain images. *Clin Radiol*. 78:e13–e21 2023.
2. Akai H, Yasaka K, Sugawara H, Tajima T, Kamitani M, Furuta T, Akahane M, Yoshioka N, Ohtomo K, Abe O, Kiryu S Acceleration of knee magnetic resonance imaging using a combination of compressed sensing and commercially available deep learning reconstruction: a preliminary study. *BMC Med Imaging*. 23:5 2023.
3. Sugawara H, Yatabe Y, Watanabe H, Akai H, Abe O, Watanabe SI, Kusumoto M. Radiological precursor lesions of lung squamous cell carcinoma: Early progression patterns and divergent volume doubling time between hilar and peripheral zones. *Lung Cancer*. 176:31–37 2023.
4. Ohgami Y, Kotani Y, Yoshida N, Akai H, Kunimatsu A, Kiryu S, Inoue Y. The contralateral effects of anticipated stimuli on brain activity measured by ERP and fMRI. *Psychophysiology*. 60:e14189 2023.
5. Kiryu S, Akai H, Yasaka K, Tajima T, Kunimatsu A, Yoshioka N, Akahane M, Abe O, Ohtomo K. Clinical Impact of Deep Learning Reconstruction in MRI. *Radiographics*. 43:e220133 2023.
6. Akai H, Yasaka K, Sugawara H, Tajima T, Akahane M, Yoshioka N, Ohtomo K, Abe O, Kiryu S. Commercially Available Deep-learning-reconstruction of MR Imaging of the Knee at 1.5T Has Higher Im-

- age Quality Than Conventionally-reconstructed Imaging at 3T: A Normal Volunteer Study. *Magn Reson Med Sci.* 22:353-360 2023.
7. Sugawara H, Furuta T, Sumiyoshi A, Iiyama M, Kamitani M, Suzuki A, Murakami A, Abe O, Aoki I, Akai H. Feasibility study of direct CT lymphangiography in mice: comparison with interstitial CT/MR lymphangiography. *Eur Radiol.* 33:5028-5036 2023.
 8. Hosoi R, Yasaka K, Mizuki M, Yamaguchi H, Miyo R, Hamada A, Abe O. Deep learning reconstruction with single-energy metal artifact reduction in pelvic computed tomography for patients with metal hip prostheses. *Jpn J Radiol.* 41:863-871 2023.
 9. Yamaguchi H, Ichikawa T, Morisaka H, Akai H, Izuka K, Ueno T, Abe O, Tsushima Y. Early detection of hypervascularization in hepatocellular carcinoma (≤ 2 cm) on hepatic arterial phase with virtual monochromatic imaging: Comparison with low-tube voltage CT. *Medicine (Baltimore).* 102:e34774 2023.
 10. Tajima T, Akai H, Yasaka K, Kunimatsu A, Yoshio-ka N, Akahane M, Ohtomo K, Abe O, Kiryu S. Comparison of 1.5 T and 3 T magnetic resonance angiography for detecting cerebral aneurysms using deep learning-based computer-assisted detection software. *Neuroradiology.* 65:1473-1482 2023.
 11. Sugawara H, Kikkawa N, Ito K, Watanabe H, Kaku S, Akai H, Abe O, Watanabe S, Yatabe Y, Kusumoto M. Is 18F-fluorodeoxyglucose PET recommended for small lung nodules? CT findings of 18F-fluorodeoxyglucose non-avid lung cancer. *Br J Radiol* 2023 Dec 13. [Epub ahead of print]